FIG. 1 (PRIOR ART)




FIG. 2




FIG. 3




FIG. 4




FIG. 5

START

OBTAIN m x n FILTERED IMAGES USING GARBOR FILTERS HAVING COMBINATIONS OF m SCALES AND n ORIENTATIONS WITH RESPECT TO TWO ARBITRARY IMAGES, ASSUMING THAT m AND n ARE PREDETERMINED POSITIVE INTEGERS

一502

CALCULATE MATCHING METRIC DEFINED BY EQUATION,

 $d\left(i,j\right) = \sum_{\text{m,n}} d_{\text{m,n}}\left(i,j\right) + b$, with Respect to Original images and m x n filtered images, assuming that means and variances of Pixel values of Original images and the respective images are denoted by μ and σ , and μ_0 and σ_0 , respectively,

$$b = \left| \frac{\mu_0^{(i)} - \mu_0^{(j)}}{\alpha(\mu_0)} \right| + \left| \frac{\sigma_0^{(i)} - \sigma_0^{(j)}}{\alpha(\sigma_0)} \right| ,$$

$$d_{\mathsf{m,n}}(i,j) = \left| \frac{\mu_{\mathsf{m,n}}^{(i)} - \mu_{\mathsf{m,n}}^{(j)}}{\alpha(\mu_{\mathsf{m,n}})} \right| + \left| \frac{\sigma_{\mathsf{m,n}}^{(i)} - \sigma_{\mathsf{m,n}}^{(j)}}{\alpha(\sigma_{\mathsf{m,n}})} \right|$$

-504

END

FIG. 6



OBTAIN m x n FILTERED IMAGES USING GARBOR FILTERS HAVING COMBINATIONS OF m SCALES AND n ORIENTATIONS WITH RESPECT TO TWO ARBITRARY IMAGES, ASSUMING THAT m AND n ARE PREDETERMINED POSITIVE INTEGERS

-602

-604

CALCULATE MATCHING METRIC DEFINED BY EQUATION,

$$d_{m,n}(i,j) = \min_{1 \le t \le K} \left[\sum_{m,n} \left(\left| \frac{\mu_{m,n}^{(i)}}{\alpha(\mu_{m,n})} - \frac{\mu_{m,n}^{(j)} \oplus t}{\alpha(\mu_{m,n})} \right| \right]$$

$$+ \left| \frac{\sigma_{\mathsf{m,n}}^{(i)}}{\alpha(\sigma_{\mathsf{m,n}})} - \frac{\sigma_{\mathsf{m,n}}^{(j)} \oplus t}{\alpha(\sigma_{\mathsf{m,n}})} \right| \right) \right] + b,$$

WITH RESPECT TO m x n FILTERED IMAGES, ASSUMING THAT MEAN AND VARIANCE OF PIXEL VALUES OF THE RESPECTIVE IMAGES ARE DENOTED BY μ AND σ , RESPECTIVELY, MEAN AND VARIANCE OF PIXEL VALUES OF ORIGINAL IMAGES ARE DENOTED BY μ_0 AND σ_0 RESPECTIVELY, K IS A PREDETERMINED POSITIVE INTEGER REPRESENTING THE NUMBER OF ORIENTATION COEFFICIENTS TO BE CONSIDERED,

$$b = \left| \frac{\mu_0^{(i)} - \mu_0^{(j)}}{\alpha(\mu_0)} \right| + \left| \frac{\sigma_0(i) - \sigma_0(j)}{\alpha(\sigma_0)} \right| + \Delta ND \quad (1)$$

DENOTES A MODULO-SHIFT FUNCTION

END

FIG. 7



OBTAIN m x n FILTERED IMAGES USING GARBOR FILTERS HAVING COMBINATIONS OF m SCALES AND n ORIENTATIONS WITH RESPECT TO TWO ARBITRARY IMAGES, ASSUMING THAT m AND n ARE PREDETERMINED POSITIVE INTEGERS

-702

-704

CALCULATE MATCHING METRIC DEFINED BY EQUATION,

$$d_{m,n}(i,j) = \min_{\substack{p=0,1\\q=0,1}} \left[\sum_{m=1}^{S-1} \sum_{m} \left(\left| \frac{\mu_{m+p,n}^{i}}{\alpha(\mu_{m+p,n})} - \frac{\mu_{m+q,n}^{j}}{\alpha(\mu_{m+q,n})} \right| \right]$$

$$+ \left| \frac{\sigma_{\mathsf{m+p,n}}^{i}}{\alpha(\sigma_{\mathsf{m+p,n}})} - \frac{\sigma_{\mathsf{m+q,n}}^{j}}{\alpha(\sigma_{\mathsf{m+q,n}})} \right| \right) \right] + b$$

WITH RESPECT TO m x n FILTERED IMAGES, ASSUMING THAT MEAN AND VARIANCE OF PIXEL VALUES OF THE RESPECTIVE IMAGES ARE DENOTED BY μ AND σ , RESPECTIVELY, MEAN AND VARIANCE OF PIXEL VALUES OF ORIGINAL IMAGES ARE DENOTED BY μ_0 AND σ_0 RESPECTIVELY, AND

$$b = \left| \frac{\mu_0^{(i)} - \mu_0^{(j)}}{\alpha(\mu_0)} \right| + \left| \frac{\sigma_0(i) - \sigma_0^{(j)}}{\alpha(\sigma_0)} \right|$$

END